

Last time: $L^p(X)$ is a Banach space

Today: Hilbert space and $L^2(X)$

Fourier transforms in general

Fix a measure space $(X, \mathcal{M}, \mu)$

Given $\psi_1, \psi_2 \in L^2(X)$ we define

$$\begin{aligned}\langle \psi_1, \psi_2 \rangle &= \int_X \psi_1^* \psi_2 \, d\mu \\ &= \int_X \psi_1^*(x) \psi_2(x) \, d\mu(x)\end{aligned}$$

By the Cauchy-Schwarz inequality

(a special case of Hölder's for $p=q=2$)

$$|\langle \psi_1, \psi_2 \rangle| \leq \int |\psi_1^* \psi_2| \, d\mu$$

$$\begin{aligned}
&= \|\psi_1^* \psi_2\|, \\
&\leq \|\psi_1^*\|_2 \|\psi_2\|_2 \\
&= \|\psi_1\|_2 \cdot \|\psi_2\|_2 < \infty
\end{aligned}$$

So $\langle \psi_1, \psi_2 \rangle$ is a well-defined complex number.

This defines an inner product on $L^2(X)$

such that $\sqrt{\langle \psi | \psi \rangle} = \|\psi\|_2$.

Since $L^2(X)$ is complete, it is a Hilbert space.

In the homework you will show that any bounded operator L on a Hilbert space $\mathcal{H}$ has an adjoint L^* , which is the unique bounded linear operator st

$$\langle L^* v, w \rangle = \langle v, L w \rangle$$

For all $v, w \in \mathcal{H}$.

So we can define normal, self-adjoint, and unitary operators as usual.

To understand the spectral theorem for Hilbert spaces, we need to generalize our notion of "diagonal operators".

We'll say a **null set** to mean a set $N \in \mathcal{M}$ st $\mu(N) = 0$.

Def: A measurable function $f: X \rightarrow \mathbb{C}$ is **essentially bounded** if $\exists$ a null set N such that f restricted to $X \setminus N$ is a bounded function.

As a result of the problem,

$$L^\infty(X) = \{ f: X \rightarrow \mathbb{C} \mid f \text{ is essentially bounded} \}.$$

If $f \in L^\infty(X)$ and $\psi \in L^2(X)$

$$\|f\psi\|_2^2 = \int_X |f\psi|^2 d\mu$$

We know $\exists$ a null set N st

$$\sup_{x \in X \setminus N} |f(x)| = \|f\|_\infty$$

If we set $\tilde{f}(x) = 0$ if $x \in N$

$f(x)$ if $x \in X \setminus N$

Then f and $\tilde{f}$ agree almost everywhere

$$\text{so } \|f\psi\|_2^2 = \int_X |\tilde{f}\psi|^2 d\mu$$

$$\leq \int_X \|f\|_\infty^2 |\psi|^2 d\mu$$

$$= \|f\|_\infty^2 \|\psi\|_2^2$$

$$\text{Hence } \|f\psi\|_2 \leq \|f\|_\infty \|\psi\|_2.$$

Given $f \in \mathcal{Z}^\infty(X)$, there is a linear operator L_f on $L^2(X)$ defined by

$$L_f(\psi) = f\psi.$$

By the inequality above, $L_f(\psi)$ is also in $L^2(X)$.

It also tells us that $\|L_f\| \leq \|f\|_\infty$.

When X is nice (σ -finite), $\|L_f\| = \|f\|_\infty$.

$$L_{f_1+f_2} = L_{f_1} + L_{f_2}$$

$$L_{\lambda f} = \lambda L_f$$

$$L_{f_1 f_2} = L_{f_1} L_{f_2}$$

So this defines an algebra map (a linear map preserving products) from

$$\mathcal{Z}^\infty(X) \rightarrow \mathcal{B}(L^2(X))$$

If f is 0 a.e. then $L_f = 0$ exactly.

Since $f\psi$ is 0 a.e. and therefore
 $f\psi = 0$ in $L^2(X)$.

Because $L^\infty(X)$ is $Z^\infty(X) / \{f = 0 \text{ a.e.}\}$

So we get a well-defined map

$$L^\infty(X) \longrightarrow B(L^2(X)) .$$

Def: We say that a ^{bounded} operator L on $L^2(X)$ is
diagonal if $L = L_f$ for some $f \in L^\infty(X)$.

The spectral theorem says that every
normal operator on a Hilbert space
can be made diagonal after a unitary
transformation

Recall a unitary map $V \rightarrow W$ is a linear isomorphism which preserves inner products.

Thm (spectral Theorem): Let $\mathcal{H}$ be a Hilbert space and let $L \in \mathcal{B}(\mathcal{H})$ be normal.

Then there exists a measure space X and a unitary map

$$U: \mathcal{H} \rightarrow L^2(X)$$

such that ULL^{-1} is a diagonal operator on $L^2(X)$. That is, $\exists f \in L^\infty(X)$ st $ULL^{-1} = L_f$.

There other ways to state the spectral theorem using projection-valued measure (equivalently, subspace valued measures from the class on d -maps)

e.g. the function $E \mapsto \Pi_E^X$ in the Problem.

$\uparrow$
measurable set

See the scratch notes in the Drive or
Hall's book for more.

We'll just say for now that given
a normal operator L on $\mathcal{H}$ there is
a canonical spectral decomposition

$$\mathcal{H} \cong \hat{\int}_{\sigma(L)}^{\oplus} \mathcal{H}_{\lambda} d\mu(\lambda)$$

↖ direct integral
of Hilbert spaces

where $\sigma(L)$ is the spectrum of L

and μ is some measure on
the Borel sets of $\sigma(L)$.

There's also a version for commuting
sets of operators:

Thm: Let S be a set of commuting normal operators on $\mathcal{H}$.

Then we can unitarily diagonalize.

every element of S simultaneously,

i.e. $\exists$ a unitary $U: \mathcal{H} \rightarrow L^2(X)$

st $L \mapsto U L U^{-1}$ sends elements of S to diagonal operators.

If S is maximal among sets of commuting normal operators, then this gives

a bijection of S with diagonal operators

on $L^2(X)$. Furthermore, in this case, X and U are unique up to equivalence.

Let G be a locally compact topological ^{abelian} group. That is, G is

an abelian group, and a Hausdorff locally compact topological space, such

that addition is a continuous map
 $G \times G \rightarrow G$ and inversion is continuous $G \rightarrow G$.

E.g. $\mathbb{Z}/N$ or $\mathbb{Z}$ w/ discrete topology
 $\mathbb{R}$

$$U(1) = S^1 = \mathbb{R}/2\pi\mathbb{Z}$$

Such a group has a unique (up to scalar)
translation-invariant measure on its Borel
sets, satisfying a regularity condition
(Radon measure). This is the Haar measure.

E.g. Counting measure on $\mathbb{Z}/n$ and $\mathbb{Z}$

Lebesgue measure on $\mathbb{R}$

Lebesgue measure on S^1

(identifying it with $[0, 2\pi)$)

$\Rightarrow$ Canonical $L^2(G)$ for any such group.

For each $g \in G$, get a unitary operator

$$T_g: L^2(G) \rightarrow L^2(G)$$

$$\psi(x) \mapsto \psi(x+g)$$

Since G is abelian, these operators all commute

$$\text{Let } S = \text{Span}\{T_g \mid g \in G\} \subseteq B(L^2(G))$$

It turns out this is a maximal set
of commuting normal operators.

By the Spectral Theorem, there is a canonical

$$U: L^2(G) \xrightarrow{\sim} L^2(X)$$

mapping S bijectively onto $L^\infty(X) \subseteq B(L^2(X))$
 $\nwarrow$ the diagonal operators.

It turns out we can take

$$X = \widehat{G} = \left\{ \chi: G \rightarrow \mathbb{C} \mid \chi \text{ is a continuous group hom.} \right\}$$

and $U T_g U^{-1}$ sends $\tilde{\psi} \in L^2(\widehat{G})$

to the function $\chi \mapsto \chi(g) \tilde{\psi}(\chi)$

The map U sends $\psi \in L^2(G)$
to its Fourier transform $\tilde{\psi} = U\psi$
 $\in L^2(\widehat{G})$

Exercise: Determine $\widehat{\mathbb{Z}}$ and $\widehat{\mathbb{R}}$.

If $\psi \in L^2(G) \cap L^1(G)$ then we can compute

$$\tilde{\psi}(\chi) = \int_G \overline{\chi(g)} \psi(g) d\mu(g).$$